

POSTOPERATIVE INFORMATION

SETH C. GAMRADT, MD

ORTHOPAEDIC SURGERY AND SPORTS MEDICINE Keck School of Medicine of USC 1520 San Pablo Street, Suite 2000 Los Angeles, CA 90033 Phone: 323.442.5860 Fax: 323.442.6952 www.gamradtortho.com Keck School of Medicine of USC

PATIENT STICKER

POST---OPERATIVE INSTRUCTIONS ULNAR COLLATERAL LIGAMENT RECONSTRUCTION

- 1. A plaster splint will be placed on your arm after surgery. You will not be able to bend or straighten your arm. However, gently make a loose fist and spread your fingers apart to keep the circulation moving in your arm.
- 2. Pain medication (PERCOCET) and an anti---inflammatory (NAPROSYN) are prescribed for you following surgery. Use the pain medication when needed. Use the anti---inflammatory as prescribed as it will help keep the swelling to a minimum and will help control pain.
- 3. You may shower. Cover the arm with a plastic bag and secure it around your upper arm with tape.
- 4. Your fingers may become swollen. Moving them frequently and elevating your arm above your heart will help minimize this.
- 5. You may notice SLIGHT numbness/tingling in your fingers. This is normal for the first 12---24 hours. If this persists or worsens, contact the office. You may be directed to loosen the ace bandage on your arm.
- 6. You may use your hand to write. You may NOT lift or push anything. Because the elbow is nearly straight, it may be difficult to use it to eat, etc.
- 7. Because the splint is heavy, your shoulder may become sore. Shrug your shoulders frequently to help minimize this.
- 8. Please call the office to schedule a follow---up appointment for wound check and application of brace, 10 days post---operatively.

IF YOU HAVE ANY QUESTIONS, PLEASE FEEL FREE TO CALL OUR OFFICE.