

POSTOPERATIVE INFORMATION

SETH C. GAMRADT, MD

ORTHOPAEDIC SURGERY AND SPORTS MEDICINE Keck School of Medicine of USC 1520 San Pablo Street, Suite 2000 Los Angeles, CA 90033

Phone: 323.442.5860 Fax: 323.442.6952

www.gamradtortho.com



PATIENT STICKER

POST---OPERATIVE INSTRUCTIONS – ARTHROSCOPIC ROTATOR CUFF REPAIR

WOUND C	ARE				
	You may remove the Operative Dressing on PostOp Day #2. THERE ARE WHITE TAPES DIRECTLY ON THE SKIN. THESE ARE STERISTRIPS—THESE SHOULD REMAIN ON.				
	Apply BandAids to the wounds. Change the BandAids daily until wounds are completely dry. Please do not use Bacitracin or other ointments under the bandage.				
	Use Ice or cold therapy device 3040 minutes 34 times per day until swelling is down, then as needed for pain relief.				
	You may shower on PostOp Day #2. Please cover the wound with plastic wrapand secure it to your skin with tape. You may remove the sling for showering, but keep the arm across the chest. Gently pat the area dry. <u>Do not soak</u> the shoulder in water. <u>Do not go swimming</u> in the pool or ocean until clear by Dr. Gamradt.				
EXERCISES					
	Wear the sling at all times except when doing your xercises. You may remove the sling for showering, but keep the arm across the chest. The sling will remain on for 46 weeks. Sleep with the sling loosely on in a position comfortable to you.				
	Please perform the exercises on the attached Home Exercise Program 2 times daily IF INSTRUCTED BY DR. GAMRADT (for larger tears, range of motion may bedelayed 24 weeks):				
	☐ Shoulder range of motion (forward elevation 90, External Rotation0)				
	☐ Elbow / Hand / Wrist Range of Motion Exercises				
	☐ Grip Strengthening				
	☐ Scapular Tightening				
POSTOP					
	You will be given a prescriptions for pain relief for you to use postoperatively:				
	 PERCOCET, a strong narcotic. Do not take Advil, Alleve or other NSAIDS until 6 weeks postoperatively as they can inhibit healing. 				
	If you have any adverse effects with the medications, please call our office.				
	If you develop a Fever (101.5), Redness or Drainage from the surgical incision site, please call our office to arrange for an evaluation.				
FOLLOW	UP				
	Please call the office to schedule a followup appointment for your wound check, 710 days post operatively. Dr. Gamradt will prescribe physical therapy atthis time.				

IF YOU HAVE ANY QUESTIONS, PLEASE FEEL FREE TO CALL OUR OFFICE.