



POSTOPERATIVE INFORMATION

SETH C. GAMRADT, MD

ORTHOPAEDIC SURGERY AND SPORTS MEDICINE

Keck School of Medicine of USC

1520 San Pablo Street, Suite 2000

Los Angeles, CA 90033

Phone: 323.442.5860

Fax: 323.442.6952

www.gamradtortho.com

The logo for the Keck School of Medicine of USC is a red rectangular box containing the text "Keck School of Medicine of USC" in a white serif font.

Keck School of
Medicine of USC

A rectangular box with a thin black border, containing the text "PATIENT STICKER" in a black sans-serif font.

PATIENT STICKER

POST---OPERATIVE INSTRUCTIONS – SHOULDER REPLACEMENT

GENERAL INFORMATION:

1. A cryocuff is a cold therapy device that is applied postoperatively. This is used most of the first 2 days in the hospital. This will help reduce the swelling and minimize the pain.
2. Your arm will be in a sling following surgery. You will be in this sling for the next 5---6 weeks. This includes sleeping with it on as well. The most important factor about sling usage and position is comfort for you. Adjust the sling so that it is most comfortable to you. I will let you know the exact duration of sling usage at your follow---up visit.
3. You will meet a Physical Therapist in the hospital the next day. They will instruct you in some exercises that you will begin doing right away. They will also instruct you how to dress yourself and apply the sling. These exercises should be done twice daily. Avoid lifting objects with the arm. You may write and type, but not drive. Driving will commence at 5---6 weeks when the sling is removed.
4. You may be more comfortable sleeping in a semi---seated position the first few nights following surgery. Keep a pillow propped under the elbow and forearm for comfort. If you have a recliner type of chair it might be beneficial. If not that is fine too, but it would be helpful to sleep propped up with pillows behind your operated shoulder as well under your elbow and forearm. This will reduce pulling on the suture lines.
5. Prescriptions for pain medication and an anti---inflammatory will be prescribed for you. You should avoid taking pain medications on an empty stomach as it will make you nauseous. Please take a full, coated aspirin (325mg) daily for 10 days to reduce the chance of blood clots.

DISCHARGE UNTIL POST---OP VISIT:

1. Ice the shoulder using the cryocuff for 20---30 minutes at least 4 times per day for the first few days. Then use it 2---3 times a day after doing your exercises.

-
2. The bandages will be changed prior to your discharge. If there is still slight drainage from the wound when you leave the hospital, change the dressing with clean, sterile gauze daily. When the dressing is

POSTOPERATIVE INFORMATION

dry, you may change it every 2---3 days. **Do Not** remove the white steri strip tape that covers that incision. Do not put anything including bacitracin or ointments on or near the incisions.

3. You may shower 96 hours after surgery but **Do Not** get the sutures wet. Wrap your shoulder with plastic wrap and secure it with tape. Dr. Gamradt will likely clear you to get the wound wet at the wound check visit 7---10 hours postoperatively.
4. You can remove your sling for short periods only when you are sitting in a chair not doing any activity. However, your elbow must be supported by pillows. You must wear the sling if you are doing any activity, especially while you are in public places for the next 5---6 weeks.
5. Formal physical therapy will begin after 2---3 weeks. Dr. Gamradt will provide a detailed prescription for therapy when you come for your first postoperative check.
6. Do not drink alcoholic beverages or take illicit drugs when taking pain medications.
7. Return to work is determined on a case by case basis.
8. When dressing, put your operative arm in the sleeve first. When getting undressed, take your operative arm out last. Loose fitting, button---down shirts are recommended.
9. In most states it is against the law to drive while your arm is in a sling.
10. First postoperative appointment is 7---10 days post surgery. An X---ray will be taken prior to this visit and the wound will be checked, steri---strips changed.